
Supplementary Table 2: Quality of life changes during cabazitaxel therapy

EORTC QLQ-C30		Baseline	Post- treatment	P for change
		Mean (SD)	best Mean (SD)	
GLOBAL	GLOBAL	48.80	71.42	0.015*
HEALTH	HEALTH	(26.97)	(18.54)	
STATUS/QOL	STATUS/QOL			
FUNCTION	Physical	54.28	61.90	0.321
SCALES	functioning	(29.42)	(32.59)	
	Role	76.19	71.43	0.673
	functioning	(26.97)	(39.34)	
	Cognitive	64.28	80.95	0.216
	functioning	(27.93)	(22.41)	
	Social	73.80	80.95	0.629
	functioning	(30.21)	(27.93)	
	Emotional	60.71	80.95	0.170
	functioning	(30.69)	(24.86)	
SYMPTOM/ITEM	Fatigue	45.23	42.85	0.853
SCALES		(31.49)	(35.81)	
	Nausea	28.57	16.66	0.140
	Vomiting	(24.93)	(30.42)	
	Pain	47.61	35.71	0.334

	(36.55)	(45.57)	
Dyspnoea	9.52 (16.26)	9.52 (16.26)	1
Insomnia	47.61(37.79)	28.57 (35.63)	0.321
Appetite	47.61 (42.41)	28.57 (40.49)	0.231
Constipation	71.42 (40.49)	19.0(37.79)	0.017*
Diarrhoea	23.80(37.08)	14.28 (26.22)	0.172
Financial difficulties	33.33 (47.14)	14.29 (37.79)	0.231
<hr/>			
EORTC QLQ-PR25	Baseline	Post- treatment	P for change
	Mean (SD)	best Mean (SD)	
<hr/>			
'Urinary symptoms	38.69 (19.5)	20.23 (24.58)	0.247
Bowel symptoms	14.28 (16.46)	5.95 (7.92)	0.234
Treatment related symptoms	23.01 (17.39)	7.93 (12.77)	0.170
Sex Interest	4.76 (12.59)	4.76 (12.59)	1.0
<hr/>			

*p<0.05, significant

Abbreviations: QOL – Quality of life; EORTC QLQ-PR25 - European Organisation for Research and Treatment of Cancer QOL Questionnaire-Prostate Module; EORTC QLQ-C30 - European Organisation for Research and Treatment of Cancer (EORTC) core QOLquestionnaire
