
## **Methotrexate Induces Apoptosis of Post-partum Placental Cytotrophoblast**

HaoYan<sup>1</sup>, Zhihua Li<sup>1</sup>, Zhonghai Yan<sup>2</sup>, Xiuying Li<sup>1</sup>, Xiaomei Li<sup>1</sup>, Lin Yu<sup>1</sup>, Chunhong Su<sup>1</sup>, Dunjin Chen<sup>1</sup>

<sup>1</sup>Department of Obstetrics and Gynecology, The Third Affiliated Hospital of Guangzhou Medical University, Key Laboratory for Major Obstetric Diseases of Guangdong Province, Guangzhou, China

<sup>2</sup>Division of Nephrology, College of Physicians and Surgeons, Columbia University, New York, NY, United States of America



## S2 Fig. Tunel index of control groups and groups with MTX treatment

First, second trimester and post-partum placental cytotrophoblast were treated with four different concentrations of MTX ( $10^{-2}$ to  $10^{-5}$  mol/L) for 48 hours. *TUNEL*, *Terminal deoxynucleotidyl transferase-mediated dUTP nick end labeling; MTX*, *methotrexate; FTC, first trimester cytotrophoblast; STC, Second trimester cytotrophoblast; PPTC: Post-partum placental cytotrophoblast.*