

Supplementary Figure 1:

Definition of the PACE brief limited intermittent psychotic symptoms (BLIPS) criteria over time



Supplementary Figure 2:

Definition of the PACE ,trait-state' risk factors / vulnerability criteria over time





Change in functional decrease definition

Yung et al. 2004 [12] CAARMS, version 01/2002 (assessment period and age range not specified) 30% drop in GAF score instead of 30 points on GAF score from premorbid level.



•Change definition of functional decline from GAF to SOFAS •Change in restriction of maximum and minimum age

Yung et al., CAARMS, version 12/2006 [13] (restriction in age range since 2005: age 15-25 yrs.) Family history of psychosis in 1st-degree relative OR Schizotypal Personality Disorder in identified patient *PLUS*

30% drop in SOFAS score from premorbid level, sustained for 1 month, occurred within past 12 months OR

SOFAS score of 50 or less for past 12 months or longer

- BDI: Beck Depression Inventory (Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J: An inventory for measuring depression. Arch Gen Psychiatry 1961;4:561-571.)
- EWSS: Early Warning Signs Survey (Birchwood M, Smith J, Macmillan F, Hogg B, Prasad R, Harvey C, Bering S: Predicting relapse in schizophrenia: the development and implementation of an early signs monitoring system using patients and families as observers, a preliminary investigation. Psychol Med 1989;19(3):649-56.)