

Supplementary Table 1. Multivariable analyses for 7-day NIHSS and 3-month mRS scores when excluding posterior circulation stroke (n=51)

	7-day NIHSS, categorized	3-month mRS	
		Adjusted OR [95% CI] p value	Adjusted OR [95% CI] p value
Model 1†			
Initial NIHSS, point, per 1 point increase	1.14 [1.03-1.27]	0.011	1.08 [0.99-1.19] 0.091
SBP, mmHg, per 10 increase	1.42 [1.12-1.80]	0.004	1.28 [1.04-1.57] 0.021
Symptom onset to IA thrombolysis, min, per 10 increase	0.94 [0.81-1.08]	0.347	0.94 [0.83-1.07] 0.349
FLAIR-hyperintense lesion	3.70 [1.13-12.10]	0.031	1.71 [0.60-4.93] 0.318
Model 2‡			
DWI lesion volume, mL, per 10 increase	1.03 [1.01-1.05]	0.001	1.02 [1.01-1.04] 0.004
SBP, mmHg, per 10 increase	1.58 [1.22-2.05]	0.001	1.33 [1.08-1.65] 0.008
Symptom onset to IA thrombolysis, min, per 10 increase	0.92 [0.80-1.06]	0.228	0.94 [0.83-1.07] 0.350
FLAIR-hyperintense lesion	3.08 [0.87-10.85]	0.081	1.40 [0.47-4.19] 0.545

Abbreviations: NIHSS, National Institutes of Health Stroke Scale; mRS, modified Rankin scale; OR, Odd ratio; CI, Confidence interval; SBP, systolic blood pressure; IA, intra-arterial; FLAIR, fluid-attenuated inversion recovery.

[†]Model 1: Ordinal logistic regression analysis of categorized 7-day NIHSS and 3-month mRS scores, adjusted for initial NIHSS score, systolic blood pressure, interval from onset to IA thrombolysis, and FLAIR-hyperintense lesion. ($p < 0.20$ for the FHL-negative and FHL-positive groups.)

[‡]Model 2: Ordinal logistic regression analysis of categorized 7-day NIHSS and 3-month mRS scores, adjusted for DWI lesion volume, systolic blood pressure, interval from onset to IA thrombolysis, and FLAIR-hyperintense lesion. ($p < 0.20$ for the FHL-negative and FHL-positive groups.)