

Novel Insights from Clinical Practice

Mismatch Repair Deficiency Drives Durable Complete Remission by Targeting Programmed Death Receptor 1 in a Metastatic Luminal Breast Cancer Patient

Carlo Fremd^a Mario Hlevnjak^b Marc Zapatka^b Inka Zoernig^a Niels Halama^a Nino Fejzibegovic^a Verena Thewes^b Peter Lichter^b Peter Schirmacher^c Matthias Kloor^c Frederik Marmé^d Florian Schütz^d Zeynep Kosaloglu^e Hans Peter Sinn^c Dirk Jäger^a Andreas Schneeweiss^a

^aNational Center for Tumor Diseases, Department of Medical Oncology, University of Heidelberg, Heidelberg, Germany; ^bDivision of Molecular Genetics, German Cancer Research Center, Heidelberg, Germany; ^cDepartment of Applied Tumor Biology, Institute of Pathology, University of Heidelberg, Heidelberg, Germany; ^dDepartment of Obstetrics and Gynecology, University of Heidelberg, Heidelberg, Germany; ^eDepartment of Vaccine Discovery, La Jolla Institute for Allergy and Immunology, La Jolla, CA, USA

Suppl. table 1. Identified somatic mutations by whole exome sequencing

Suppl. table 2. Predicted neoantigens with high-affinity binding

Please see xls files.

Suppl. fig. 1. Intermutation distances of somatic mutations.

