Supplementary Material

The Differentially Expressed Circular RNAs in the Substantia Nigra and Corpus Striatum of Nrf2-Knockout Mice

Jun-Hui Yang^a Run-Jiao Zhang^b Jia-Jie Lin^c Ming-Chao Cao^d Qie Wang^b Hui-Xian Cui^b Shao-Guang Sun^c Lei Wang^b

^aDepartment of Neurology, Hospital of Jingxing county, ^bDepartment of Human Anatomy, Hebei Medical University, ^cDepartment of Biochemistry and Molecular Biology, Hebei Medical University, ^dDepartment of rehabilitation, The second hospital of Hebei Medical University




Figure S1. DEcircRNA-miRNA-DEceRNA interaction subnetworks of mmu_circRNA_015216, mmu_circRNA_32463 and mmu_circRNA_34132 in Nrf2 (-/-)substantia nigra and DEcircRNA-miRNA-DEceRNA interaction subnetworks of mmu_circRNA_017077 and mmu_circRNA_34132 in Nrf2 (-/-)corpus striatum.

(A). Subnetwork of mmu_circRNA_015216 in Nrf2 (-/-)substantia nigra; (B). Subnetwork of mmu_circRNA_32463 in Nrf2 (-/-)substantia nigra; (C). Subnetwork of mmu_circRNA_34132 in Nrf2 (-/-) substantia nigra; (D). Subnetwork of mmu_circRNA_017077 in Nrf2 (-/-) corpus striatum; (E). Subnetwork of mmu_circRNA_34132 in Nrf2 (-/-) corpus striatum.

Brown nodes indicated DEcircRNAs. Red and blue nodes indicated miRNAs targeted by DEcircRNAs and the gene ID of their DEceRNAs, respectively. Edges represented interactions.