

Supplemental Table 1. Characteristics of participants excluded from analysis who entered CKD stage 5 in CRIC.

Baseline Characteristics	Included for analysis (N=772) N (%) or Mean $\pm$ SD	Excluded from analysis (N=98)	P-value for difference
Median age (years) [interquartile range]	60.4 [51.0-67.3]	66.4 [57.3-71.7]	<0.001
Female	330 (42.8)	50 (51)	0.12
Race/ethnicity			0.52
White	189 (24.5)	24 (25)	
Black	407 (52.7)	46 (47)	
Hispanic	142 (18.4)	24 (24)	
Obese	442 (57.3)	54 (55)	0.68
Any cardiovascular disease	344 (44.6)	49 (50)	0.31
Diabetes	512 (66.3)	61 (62)	0.42
Systolic BP (mm Hg) [IQR]	138.7 [122.7-156.7]	132.7 [120.7-148.7]	0.08
Current smoker	120 (15.5)	17 (17)	0.21
Serum potassium meq/L	4.5 $\pm$ 0.6	4.5 $\pm$ 0.6	0.83
Hemoglobin (g/dL)	11.4 $\pm$ 1.6	11.4 $\pm$ 1.4	0.62
Serum CO ₂ (mmol/L)	22.1 $\pm$ 3.5	21.6 $\pm$ 3.9	0.06
Serum Urea Nitrogen (mg/dL)	50.0 $\pm$ 17.5	57 $\pm$ 20.1	<0.001
Serum albumin (g/dL)	3.7 $\pm$ 0.5	3.7 $\pm$ 0.4	0.15
Urine protein/creatinine ratio (g/g) [IQR]	1.9 [0.75-4.0]	1.1 [0.5-2.5]	<0.001