

## Supplementary Material

### Dimethylarginine Dimethyl Aminohydrolase-1 polymorphisms and venous intimal hyperplasia in dialysis patients

Wu C.C., Hsieh M.Y., Lee C.K., Chuang S.Y., Chung M.Y., Lin C.C.

**Supplemental Table 2** Distribution of lesion locations stratified by types of vascular access

Variables	Total cohort	No early restenosis	Early restenosis	P Value
<b>Graft</b>	190	99	91	
Graft-vein junction	83 (44%)	46 (47%)	37 (41%)	0.47
Graft-artery junction	10 (5%)	6 (6%)	4 (4%)	0.75
Intra-graft	33 (17%)	14 (14%)	19 (21%)	0.25
Cephalic vein	30 (16%)	16 (16%)	14 (15%)	0.99
Basilic vein	60 (32%)	25 (26%)	35 (39%)	0.08
Others	4 (2%)	2 (2%)	2 (2%)	0.99
<b>Fistula</b>	283	201	82	
Proximal vein	158 (56%)	125 (62%)	33 (40%)	<0.01
Cephalic vein	76 (27%)	46 (23%)	30 (37%)	0.03
Basilic vein	41 (15%)	25 (13%)	16 (20%)	0.19
Others	12 (4%)	9 (4%)	3 (4%)	0.76

Data are expressed as N (%); some vessels had more than one lesion treated.