

Supplemental Table 1. Regular medications in the study groups.

	No-CV-medication (n=70)	Hypertension (n=35)	CKD 5 or 5D (n=35)
ACE inhibitor	0	9	7
AT receptor blocker	0	16	15
Beta blocker	0	14	24
Beta and alfa blocker	0	0	1
Calcium channel blocker	0	21	24
Thazide diuretic	0	7	3
Furosemide	0	0	17
Potassium sparing diuretic	0	4	0
Nitrate	0	0	1
Prazocin	0	3	3
Moxonidin	0	2	2
Statins	0	13	20
Ezetimib	0	0	0
Low dose ASA	0	5	11
DPP4 inhibitors	0	1	1
Insulins	0	1	7
Metformin	0	3	0
Female hormones	2	0	1
Thyroxin	0	0	1
Protone pump inhibitors	2	2	5
Mental health medications	2	3	4
Pregabalin	1	0	2
Alphacalcidol or paricalcitol	0	0	12
Allopurinol	1	0	7
Calcium carbonate	0	0	31
Cyclosporine A	0	0	1
Tacrolimus	0	0	2
Corticosteroid, per oral	0	0	2
Erythropoietin	0	0	23
Sodium bicarbonate	0	0	12
Inhaled corticosteroid	1	0	0
Phosphate binders (non-calcium)	0	0	6
Azathioprine	0	0	1
Methotrexate	0	1	0
Hydroxychloriquine	0	1	0
Denosumab	0	0	1
Antihistamines	1	2	0
Acyclovir	1	0	0
Pramipexol	0	0	1
Warfarin	0	1	1