

Table e1. Three-month mortality, good functional outcomes and serious cardiac adverse events according to low, moderate and high pre-stroke CHADS₂ and CHA₂DS₂-VASc scores and atrial fibrillation

	AF n=1,755	No AF n=4,857	P
Ninety-day mortality, % (95% CI)			
CHADS ₂ score			
Low risk (0)	12.7 (7.3-20.1)	9.1 (7.1-11.4)	0.2
Intermediate risk (1)	19.4 (15.6-23.6)	12.1 (10.5-13.9)	<0.001
High risk (2-6)	26.7 (24.2-29.2)	18.7 (17.3-20.3)	<0.001
CHA ₂ DS ₂ -VASc score			
Low risk (0)	7.7 (0.9-25.1)	6.6 (4.1-9.9)	0.7
Intermediate risk (1)	11.1 (6.2-17.9)	8.3 (6.5-10.4)	0.3
High risk (2-9)	25.4 (23.3-27.6)	17.6 (16.4-18.8)	<0.001
Ninety-day good functional outcomes (mRS ≤ 1), % (95% CI)			
CHADS ₂ score			
Low risk (0)	30.6 (21.7-40.7)	37.3 (33.8-41.0)	0.2
Intermediate risk (1)	27.5 (22.9-32.6)	36.4 (33.8-39.1)	0.002
High risk (2-6)	17.4 (15.2-19.7)	24.2 (22.6-26.0)	<0.001
CHA ₂ DS ₂ -VASc score			
Low risk (0)	23.5 (6.8-49.9)	38.8 (33.4-44.5)	0.3
Intermediate risk (1)	37.6 (28.5-47.4)	42.4 (38.8-46.0)	0.4
High risk (2-9)	19.1 (17.1-21.2)	26.4 (24.9-27.9)	<0.001
Ninety-day serious cardiac adverse events, % (95% CI)			
CHADS ₂ score			
Low risk (0)	11.0 (6.0-18.1)	4.7 (3.3-6.5)	0.02
Intermediate risk (1)	12.6 (9.5-16.3)	6.6 (5.4-6.0)	<0.001

High risk (2-6)	14.5 (12.6-16.6)	9.5 (8.5-10.7)	<0.001
CHA ₂ DS ₂ -VASc score			
Low risk (0)	3.8 (0.1-19.6)	3.8 (2.0-6.5)	1.0
Intermediate risk (1)	7.9 (3.9-14.1)	4.2 (2.9-5.8)	0.07
High risk (2-9)	14.5 (12.8-16.3)	9.1 (8.2-10.1)	<0.001

Abbreviations: CI = interquartile range.